



## The canonical and alternate duals of a wavelet frame

Lemvig, Jakob

*Publication date:*  
2007

[Link back to DTU Orbit](#)

*Citation (APA):*  
Lemvig, J. (Author). (2007). The canonical and alternate duals of a wavelet frame. Sound/Visual production (digital)

---

### General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

# The Canonical and Alternate Duals of a Wavelet Frame

Jakob Lemvig

[Joint with: Marcin Bownik, University of Oregon]

Department of Mathematics, Technical University of Denmark (DTU)

Summer School, Inzell  
September 17, 2007

## Motivation.

Wavelet frames: A generalization of wavelet orthonormal basis with more flexibility and freedom.

## Motivation.

Wavelet frames: A generalization of wavelet orthonormal basis with more flexibility and freedom.

Do we always have this freedom?

## Motivation.

Wavelet frames: A generalization of wavelet orthonormal basis with more flexibility and freedom.

Do we always have this freedom?

The natural representation of a function in terms of a frame involves the so-called canonical dual. The canonical dual of a wavelet frame need not have wavelet structure; and even worse, there might **not be any** dual with wavelet structure.

## One Result in this Direction

Suppose  $\psi$  is an orthonormal wavelet and let

$$\theta(x) = \psi(x) + \varepsilon 2^{1/2} \psi(2x)$$

for  $0 < \varepsilon < 1$ . Then  $\theta$  generates a wavelet Riesz basis whose (canonical) dual is **not** of the form

$$\{2^{j/2} \phi(2^j x - k) : j, k \in \mathbb{Z}, \phi \in \Phi\}$$

for any finite set  $\Phi$  of generators ([Daubechies], [Chui & Shi]).

## One Result in this Direction

Suppose  $\psi$  is an orthonormal wavelet and let

$$\theta(x) = \psi(x) + \varepsilon 2^{1/2} \psi(2x)$$

for  $0 < \varepsilon < 1$ . Then  $\theta$  generates a wavelet Riesz basis whose (canonical) dual is **not** of the form

$$\{2^{j/2} \phi(2^j x - k) : j, k \in \mathbb{Z}, \phi \in \Phi\}$$

for any finite set  $\Phi$  of generators ([Daubechies], [Chui & Shi]). Obviously, the Riesz wavelet has no alternate wavelet duals either, so one might ask:

*Does the existence of a dual wavelet frame imply wavelet structure of the canonical dual?*

## Main result of this work

In this talk we will explore the relationship between canonical and alternate dual frames of a wavelet frame. The main result is a **negative answer** to the question on the previous slide:

### Theorem (Bownik, L.)

*For all  $J \in \mathbb{N}$ , there exists a frame wavelet  $\psi \in L^2(\mathbb{R})$  such that:*

- (i)  $\hat{\psi}$  is  $C^\infty$  and compactly supported,*
- (ii) its canonical dual frame is not a wavelet system generated by fewer than  $2^J$  functions,*
- (iii) there are infinitely many  $\tilde{\psi}$  such that  $\psi$  and  $\tilde{\psi}$  form a pair of dual wavelet frames.*



## Main result of this work

In this talk we will explore the relationship between canonical and alternate dual frames of a wavelet frame. The main result is a **negative answer** to the question on the previous slide:

### Theorem (Bownik, L.)

*For all  $J \in \mathbb{N}$ , there exists a frame wavelet  $\psi \in L^2(\mathbb{R})$  such that:*

- (i)  $\hat{\psi}$  is  $C^\infty$  and compactly supported,*
- (ii) its canonical dual frame is not a wavelet system generated by fewer than  $2^J$  functions,*
- (iii) there are infinitely many  $\tilde{\psi}$  such that  $\psi$  and  $\tilde{\psi}$  form a pair of dual wavelet frames.*

This claim (with  $J = 1$ ) was asserted by Daubechies and Han [Appl. Comp. Harmonic Anal. **12** (2002), no. 3, 269–285].

## What is a Frame?

A sequence  $\{f_j\}_{j \in \mathbb{N}}$  is a **frame** for a separable Hilbert space  $\mathcal{H}$  if

$$\exists C_1, C_2 > 0 : \quad C_1 \|f\|^2 \leq \sum_{j \in \mathbb{N}} |\langle f, f_j \rangle|^2 \leq C_2 \|f\|^2 \quad \text{for all } f \in \mathcal{H}.$$

If the upper bound holds in the above inequality, then  $\{f_j\}$  is said to be a **Bessel sequence**. Two Bessel sequences  $\{f_j\}$  and  $\{g_j\}$  are said to be **dual frames** if

$$f = \sum_{j \in \mathbb{N}} \langle f, g_j \rangle f_j \quad \text{for all } f \in \mathcal{H}.$$

## What is a Frame?

A sequence  $\{f_j\}_{j \in \mathbb{N}}$  is a **frame** for a separable Hilbert space  $\mathcal{H}$  if

$$\exists C_1, C_2 > 0 : \quad C_1 \|f\|^2 \leq \sum_{j \in \mathbb{N}} |\langle f, f_j \rangle|^2 \leq C_2 \|f\|^2 \quad \text{for all } f \in \mathcal{H}.$$

If the upper bound holds in the above inequality, then  $\{f_j\}$  is said to be a **Bessel sequence**. Two Bessel sequences  $\{f_j\}$  and  $\{g_j\}$  are said to be **dual frames** if

$$f = \sum_{j \in \mathbb{N}} \langle f, S^{-1} f_j \rangle f_j \quad \text{for all } f \in \mathcal{H}.$$

At least one dual always exists: the **canonical dual**  $\{S^{-1} f_j\}$ , where the frame operator of  $\{f_j\}$  is  $S: \mathcal{H} \rightarrow \mathcal{H}, S f = \sum \langle f, f_j \rangle f_j$ .

Redundant frames have several duals; a dual which is not the canonical dual is called an **alternate** dual.

## What is a Frame Wavelet?

The **wavelet system** generated by  $\Psi = \{\psi_1, \dots, \psi_L\}$  is defined as

$\{\psi_{j,k} : j, k \in \mathbb{Z}, \psi \in \Psi\}$ , where  $\psi_{j,k} := \psi(a^j \cdot -k) \equiv D_a^j T_k \psi \equiv \mathcal{U}(\Psi)$ .

where the **translation and dilation** operators on  $L^2(\mathbb{R})$  are:

$$T_k f(x) := f(x - k), \quad k \in \mathbb{Z},$$

$$D_a f(x) := a^{1/2} f(ax), \quad a > 1.$$

The wavelet unitaries are denoted  $\mathcal{U} := \{D_a^j T_k : j \in \mathbb{Z}, k \in \mathbb{Z}\}$ .

## What is a Frame Wavelet?

The wavelet system generated by  $\Psi = \{\psi_1, \dots, \psi_L\}$  is defined as

$$\{\psi_{j,k} : j, k \in \mathbb{Z}, \psi \in \Psi\}, \text{ where } \psi_{j,k} := \psi(a^j \cdot -k) \equiv D_a^j T_k \psi \equiv \mathcal{U}(\Psi).$$

A set of functions  $\Psi \in L^2(\mathbb{R})$  is called a **frame wavelet** if  $\{\psi_{j,k}\}_{j \in \mathbb{Z}, k \in \mathbb{Z}, \psi \in \Psi}$  is a frame for  $L^2(\mathbb{R})$ .

## What is a Frame Wavelet?

The wavelet system generated by  $\Psi = \{\psi_1, \dots, \psi_L\}$  is defined as

$\{\psi_{j,k} : j, k \in \mathbb{Z}, \psi \in \Psi\}$ , where  $\psi_{j,k} := \psi(a^j \cdot -k) \equiv D_a^j T_k \psi \equiv \mathcal{U}(\Psi)$ .

### Expansion property

Suppose  $\Psi$  is a frame wavelet. Then

$$f = \sum_{\psi \in \Psi} \sum_{j \in \mathbb{Z}} \sum_{k \in \mathbb{Z}} \langle f, S^{-1} \psi_{j,k} \rangle \psi_{j,k} \quad \text{for all } f \in L^2(\mathbb{R}),$$

## Notation

A closed subspace  $W \subset L^2(\mathbb{R})$  is said to be *M $\mathbb{Z}$  shift invariant* if  $T_{Mz}W \subset W$  for all  $z \in \mathbb{Z}$ .

Given a frame wavelet  $\Psi$ , the *space of negative dilates*  $V(\Psi)$  is:

$$V(\Psi) = \overline{\text{span}}\{\psi_{j,k} : j < 0, k \in \mathbb{Z}, \psi \in \Psi\}.$$

The *local commutant* of a system of operators  $\mathcal{A}$  at the point  $f \in L^2(\mathbb{R})$  is defined as

$$\mathcal{C}_f(\mathcal{A}) := \{B \in \mathcal{B}(L^2(\mathbb{R})) : BAf = ABf \quad \forall A \in \mathcal{A}\}.$$

# The Canonical Dual Frame

The canonical dual of a wavelet frame  $\{D_a^j T_k \psi\}_{j,k \in \mathbb{Z}, \psi \in \Psi}$  is:

$$\begin{aligned} \{S^{-1} D_a^j T_k \psi_i : j, k \in \mathbb{Z}, i = 1, \dots, L\} &= \{D_a^j S^{-1} T_k \psi_i : j, k \in \mathbb{Z}, i = 1, \dots, L\} \\ &= \{D_a^j \eta^{k,i} : j, k \in \mathbb{Z}, i = 1, \dots, L\}, \end{aligned}$$

where  $S$  is the frame operator of  $\mathcal{U}(\Psi)$ , and  $\{\eta^{k,i}\}$  is a family of functions, **not** necessarily with translation structure, indexed by  $\{1, \dots, L\} \times \mathbb{Z}$ .



## The Canonical Dual Frame

The canonical dual of a wavelet frame  $\{D_a^j T_k \psi\}_{j,k \in \mathbb{Z}, \psi \in \Psi}$  is:

$$\begin{aligned} \{S^{-1} D_a^j T_k \psi_i : j, k \in \mathbb{Z}, i = 1, \dots, L\} &= \{D_a^j S^{-1} T_k \psi_i : j, k \in \mathbb{Z}, i = 1, \dots, L\} \\ &= \{D_a^j \eta^{k,i} : j, k \in \mathbb{Z}, i = 1, \dots, L\}, \end{aligned}$$

where  $S$  is the frame operator of  $\mathcal{U}(\Psi)$ , and  $\{\eta^{k,i}\}$  is a family of functions, [not](#) necessarily with translation structure, indexed by  $\{1, \dots, L\} \times \mathbb{Z}$ . The canonical dual takes the form of a [wavelet system](#) generated by  $|\Psi| = L$  functions precisely when

$$T_k S^{-1} \psi = S^{-1} T_k \psi \quad \text{for all } \psi \in \Psi, k \in \mathbb{Z},$$

that is, precisely when  $S^{-1} \in \mathcal{C}_\psi(\{T_k : k \in \mathbb{Z}\})$  for all  $\psi \in \Psi$ .

## Lifting duality to a Sparser Lattice

For simplicity let  $L = 1$ , i.e.,  $\Psi = \{\psi\}$ . What can we “do” if  $T_k S^{-1} \psi \neq S^{-1} T_k \psi$  for some  $k \in \mathbb{Z}$ , i.e., if the canonical dual is not a wavelet system generated by one function?

## Lifting duality to a Sparser Lattice

For simplicity let  $L = 1$ , i.e.,  $\Psi = \{\psi\}$ . What can we “do” if  $T_k S^{-1} \psi \neq S^{-1} T_k \psi$  for some  $k \in \mathbb{Z}$ , i.e., if the canonical dual is not a wavelet system generated by one function?

Idea: For  $P \in \mathbb{N}$  consider  $\{D_a^j T_k \psi\}_{j,k \in \mathbb{Z}}$  as a wavelet system on the form  $\{D_a^j T_{Pk} \psi\}_{j,k \in \mathbb{Z}, \psi \in \{\psi, T_1 \psi, \dots, T_{P-1} \psi\}}$ .

## Lifting duality to a Sparser Lattice

For simplicity let  $L = 1$ , i.e.,  $\Psi = \{\psi\}$ . What can we “do” if  $T_k S^{-1} \psi \neq S^{-1} T_k \psi$  for some  $k \in \mathbb{Z}$ , i.e., if the canonical dual is not a wavelet system generated by one function?

Idea: For  $P \in \mathbb{N}$  consider  $\{D_a^j T_k \psi\}_{j,k \in \mathbb{Z}}$  as a wavelet system on the form  $\{D_a^j T_{Pk} \psi\}_{j,k \in \mathbb{Z}, \psi \in \{\psi, T_1 \psi, \dots, T_{P-1} \psi\}}$ .

Now, it might happen that *this* system has a canonical dual with wavelet structure as systems on the sparser translation lattice  $P\mathbb{Z}$  and  $P$  generators. This happens precisely when

$$T_{Pk} S^{-1} \psi = S^{-1} T_{Pk} \psi \quad \text{for all } k \in \mathbb{Z}.$$

We note that the frame operator  $S$  remains the same during this change of lattice.

## Lifting duality to a Sparser Lattice

For simplicity let  $L = 1$ , i.e.,  $\Psi = \{\psi\}$ . What can we “do” if  $T_k S^{-1} \psi \neq S^{-1} T_k \psi$  for some  $k \in \mathbb{Z}$ , i.e., if the canonical dual is not a wavelet system generated by one function?

Idea: For  $P \in \mathbb{N}$  consider  $\{D_a^j T_k \psi\}_{j,k \in \mathbb{Z}}$  as a wavelet system on the form  $\{D_a^j T_{Pk} \psi\}_{j,k \in \mathbb{Z}, \psi \in \{\psi, T_1 \psi, \dots, T_{P-1} \psi\}}$ .

Then we have  $P$  functions  $\{\phi_0, \phi_1, \dots, \phi_{P-1}\}$  such that

$$\{D^j T_{Pk}(\phi_0), D^j T_{Pk}(\phi_1), \dots, D^j T_{Pk}(\phi_{P-1})\}_{j,k \in \mathbb{Z}}$$

is the canonical dual of

$$\{D^j T_{Pk}(\psi), D^j T_{Pk}(T_1 \psi), \dots, D^j T_{Pk}(T_{P-1} \psi)\}_{j,k \in \mathbb{Z}}$$

# The Period of a Frame Wavelet

## Definition

Suppose that  $\Psi = \{\psi_1, \dots, \psi_L\} \subset L^2(\mathbb{R})$  is a frame wavelet associated with an integer dilation factor  $a$ ,  $|a| \geq 2$ . The **period** of  $\Psi$  is the smallest integer  $p \geq 1$  such that for all  $f \in \overline{\text{span}} \{T_k \psi : k \in \mathbb{Z}, \psi \in \Psi\}$ ,

$$T_{pk} S^{-1} f = S^{-1} T_{pk} f \quad \text{for all } k \in \mathbb{Z},$$

where  $S$  is the frame operator of the wavelet frame generated by  $\Psi$ . If there is no such  $p$ , we say that the period of  $\Psi$  is  $\infty$ .

## Periods and the Canonical Dual

### Proposition (Bownik, L.)

Suppose that  $\Psi = \{\psi_1, \dots, \psi_L\} \subset L^2(\mathbb{R})$  is a frame wavelet. For any nonnegative integer  $M \in \mathbb{N}$ , the following are equivalent:

- (i)  $P(\Psi) \mid M$ , i.e., the period of  $\Psi$ , denoted  $P(\Psi)$ , divides  $M$ .
- (ii) There exist  $ML$  functions  $\Phi = \{\phi_1, \dots, \phi_{ML}\}$  such that  $\{D_a^j T_{Mk} \phi\}_{j,k \in \mathbb{Z}, \phi \in \Phi}$  is the canonical dual of  $\{D_a^j T_k \psi\}_{j,k \in \mathbb{Z}, \psi \in \Psi} = \{D_a^j T_{Mk} \psi\}_{j,k \in \mathbb{Z}, \psi \in \Psi_M}$ , where

$$\Psi_M := \{T_m \psi : m = 0, \dots, M-1, \psi \in \Psi\}.$$

Hence, if the period  $P(\Psi)$  of a frame wavelet  $\Psi$  is finite, then the canonical dual frame is a wavelet system generated by  $P(\Psi) \cdot |\Psi|$  functions, and this is the least number of generators.

## Periods and the Canonical Dual

### Proposition (Bownik, L.)

Suppose that  $\Psi = \{\psi_1, \dots, \psi_L\} \subset L^2(\mathbb{R})$  is a frame wavelet. For any nonnegative integer  $M \in \mathbb{N}$ , the following are equivalent:

- (i)  $P(\Psi) \mid M$ , i.e., the period of  $\Psi$ , denoted  $P(\Psi)$ , divides  $M$ .
- (ii) There exist  $ML$  functions  $\Phi = \{\phi_1, \dots, \phi_{ML}\}$  such that  $\{D_a^j T_{Mk} \phi\}_{j,k \in \mathbb{Z}, \phi \in \Phi}$  is the canonical dual of  $\{D_a^j T_k \psi\}_{j,k \in \mathbb{Z}, \psi \in \Psi} = \{D_a^j T_{Mk} \psi\}_{j,k \in \mathbb{Z}, \psi \in \Psi_M}$ , where

$$\Psi_M := \{T_m \psi : m = 0, \dots, M-1, \psi \in \Psi\}.$$

Note: In this case the wavelet structure of the canonical dual frame is altered since it is based on the translation lattice  $P(\Psi) \cdot \mathbb{Z}$  which is sparser than the original lattice  $\mathbb{Z}$ .



## The Main Result (again)

### Theorem

*For all  $J \in \mathbb{N}$ , there exists a frame wavelet  $\psi \in L^2(\mathbb{R})$  such that:*

- (i)  $\hat{\psi}$  is  $C^\infty$  and compactly supported,*
- (ii) its canonical dual frame is not a wavelet system generated by fewer than  $2^J$  functions,*
- (iii) there are infinitely many  $\tilde{\psi}$  such that  $\psi$  and  $\tilde{\psi}$  form a pair of dual wavelet frames.*

## Idea of the Proof (case $J = 1$ )

Construct a nice frame wavelet with an alternate wavelet dual and a non  $\mathbb{Z}$  shift invariant space of negative dilates  $V(\psi)$ . Use the negation of the result below to conclude that the canonical dual cannot be generated by one function.

### Proposition (Bownik & Weber)

*Suppose that the canonical dual of a wavelet frame  $\psi$  has a wavelet structure, i.e., it is of the form  $\{\phi_{j,k} : j, k \in \mathbb{Z}\}$  for some frame wavelet  $\phi$ . Then, the space of negative dilates*

$$V(\psi) = \overline{\text{span}}\{\psi_{j,k} : j < 0, k \in \mathbb{Z}\}$$

*is shift-invariant.*

## Sketch of the Proof (general case $J \in \mathbb{N}$ )

Fix  $J \in \mathbb{N}$ . Construct a smooth frame wavelet  $\psi = \psi^0 + \varepsilon\psi^1$  as in the first lemma and the figure with  $N = J + 3$  and with a space of negative dilates  $V(\psi)$  **not** shift invariant under  $M\mathbb{Z}$  for any  $M = 1, 2, \dots, 2^J$  (second lemma).

## Sketch of the Proof (general case $J \in \mathbb{N}$ )

Fix  $J \in \mathbb{N}$ . Construct a smooth frame wavelet  $\psi = \psi^0 + \varepsilon\psi^1$  as in the first lemma and the figure with  $N = J + 3$  and with a space of negative dilates  $V(\psi)$  **not** shift invariant under  $M\mathbb{Z}$  for any  $M = 1, 2, \dots, 2^J$  (second lemma). By the result below, the period of such a framelet  $\psi$  will be at least  $2^J$ . Therefore, by the Proposition above, the canonical dual is at least generated by  $2^J$  functions. Finally, use the characteristic equations for dual wavelets frames to explicitly show that  $\psi^0$  is an alternate dual wavelet of  $\psi$ .

### Proposition (Bownik & Weber)

*Let  $M \in \mathbb{N}$ . If  $\Psi$  is a frame wavelet and the period of  $\Psi$  divides  $M$ , then  $V(\Psi)$  is shift invariant by the lattice  $M\mathbb{Z}$ .*

# First lemma: Constructing the Frame Wavelet


## Lemma

For every  $N \geq 4$  and  $0 < \delta < 2^{-N}$ , there exists a frame wavelet  $\psi$  such that  $\hat{\psi} \in C_0^\infty(\mathbb{R})$  and

$$\begin{aligned} \hat{\psi}(\xi) \neq 0 &\iff \xi \in (-1/2, -1/4) \cup (1/2, 3/4) \\ &\quad \cup \left( -2^{-N+1} - \delta, -2^{-N} + \delta \right) \\ &\quad \cup \left( 2^{-N} - \delta, 2^{-N+1} + \delta \right) \end{aligned}$$

$$\hat{\psi}(\xi) = \hat{\psi}(\xi - 1) \neq 0 \quad \text{for } \xi \in (1/2, 3/4).$$

## First lemma: Constructing the Frame Wavelet



**Figure:** Sketch of the graph of  $\hat{\psi} = \hat{\psi}^0 + \varepsilon \hat{\psi}^1$ ;  $\psi$  and  $\psi^0$  are (alternate) duals.

## Second Lemma: Showing Non Shift Invariance of $V(\psi)$

### Lemma

Suppose that  $\psi \in L^2(\mathbb{R})$  is as in the first lemma with  $N \geq 4$  and  $0 < \delta < 2^{-N}$ . Then the space of negative dilates  $V(\psi)$  is *not*  $p\mathbb{Z}$ -SI for any  $p < 2^{N-3}$ ,  $p \in \mathbb{N}$ .

### Sketch of proof.

Let  $W_j := \overline{\text{span}}\{\psi_{j,k} : k \in \mathbb{Z}, \psi \in \Psi\}$  for each  $j \in \mathbb{Z}$ . Use that  $V(\psi) = \overline{\text{span}}\bigcup_{j < 0} W_j(\psi)$ , that  $f \in W_0(\psi)$  if and only if  $f(2^j \cdot) \in W_j(\psi)$ , and that the principal shift-invariant subspace  $W_0(\psi)$  can be described as

$$W_0(\psi) = \{f \in L^2(\mathbb{R}) : \hat{f} = \hat{\psi}m \text{ for some meas., 1-periodic } m\}.$$



## Second Lemma: Showing Non Shift Invariance of $V(\psi)$

### Lemma

Suppose that  $\psi \in L^2(\mathbb{R})$  is as in the first lemma with  $N \geq 4$  and  $0 < \delta < 2^{-N}$ . Then the space of negative dilates  $V(\psi)$  is *not*  $p\mathbb{Z}$ -SI for any  $p < 2^{N-3}$ ,  $p \in \mathbb{N}$ .

### Sketch of proof.

Let  $W_j := \overline{\text{span}}\{\psi_{j,k} : k \in \mathbb{Z}, \psi \in \Psi\}$  for each  $j \in \mathbb{Z}$ . Use that  $V(\psi) = \overline{\text{span}}\bigcup_{j < 0} W_j(\psi)$ , that  $f \in W_0(\psi)$  if and only if  $f(2^j \cdot) \in W_j(\psi)$ , and that the principal shift-invariant subspace  $W_0(\psi)$  can be described as

$$W_0(\psi) = \left\{ f \in L^2(\mathbb{R}) : \begin{aligned} \text{supp } \hat{f} &\subset [-1/2, -1/4] \cup [1/2, 3/4] \cup K \\ \hat{f}(\xi - 1) &= \hat{f}(\xi) \quad \text{a.e. } \xi \in [1/2, 3/4] \end{aligned} \right\},$$

where  $K = [-2^{-N+1} - \delta, -2^{-N} + \delta] \cup [2^{-N} - \delta, 2^{-N+1} + \delta]$ .



Second Lemma: Showing Non Shift Invariance of  $V(\psi)$ 

Proof (cont.)

$$V(\psi) = \left\{ f \in L^2(\mathbb{R}) : \text{supp } \hat{f} \subset \bigcup_{j=-N+3}^{-1} 2^j([-1/2, -1/4] \cup [1/2, 3/4]), \right. \\
\hat{f}(\xi - 2^{-1}) = \hat{f}(\xi) \quad a.e. \xi \in [2^{-2}, 3/2 \cdot 2^{-2}] , \\
\hat{f}(\xi - 2^{-2}) = \hat{f}(\xi) \quad a.e. \xi \in [2^{-3}, 3/2 \cdot 2^{-3}] , \\
\vdots \\
\left. \hat{f}(\xi - 2^{-N+3}) = \hat{f}(\xi) \quad a.e. \xi \in [2^{-N+2}, 3/2 \cdot 2^{-N+2}] \right\}.$$

Define  $f \in V(\psi)$  by  $\hat{f} = \chi_{[2^{-N+2}, 3/2 \cdot 2^{-N+2}] \cup [-2^{-N+2}, -1/2 \cdot 2^{-N+2}]}$ .  
 Show that  $T_p f \notin V(\psi)$  for  $p < 2^{N-3}$ , or equivalently, that  
 $\exp(2\pi i p \cdot) \hat{f} \notin \mathcal{F}V(\psi)$ .



## In Perspective with Other Known Results

### Theorem (Aucher, 1990)

*Suppose  $\psi \in L^2(\mathbb{R})$  is an orthonormal wavelet,  $|\hat{\psi}|$  continuous, and  $\hat{\psi} = \mathcal{O}(|\xi|^{-1/2-\delta})$  for some  $\delta > 0$ . Then  $\psi$  is an MRA wavelet.*

Hence, all “regular” orthonormal wavelets are associated with an MRA.

## In Perspective with Other Known Results

### Theorem (Aucher, 1990)

*Suppose  $\psi \in L^2(\mathbb{R})$  is an orthonormal wavelet,  $|\hat{\psi}|$  continuous, and  $\hat{\psi} = \mathcal{O}(|\xi|^{-1/2-\delta})$  for some  $\delta > 0$ . Then  $\psi$  is an MRA wavelet.*

Hence, all “regular” orthonormal wavelets are associated with an MRA. Baggett et al. constructed a non-MRA  $C^r$  tight frame wavelet with rapid decay for any  $r \in \mathbb{N}$ . The non-tight frame wavelet from the main result is an example of a non-GMRA  $C^\infty$  frame wavelet with rapid decay.

## References I



M. Bownik, J. Lemvig.

The canonical and alternate duals of a wavelet frame.

*Appl. Comput. Harmon. Anal.*, **23** (2007), no. 2, 263–272.



M. Bownik, E. Weber.

Affine frames, GMRA's, and the canonical dual.

*Studia Math.*, 159(3):453–479, 2003.



I. Daubechies, B. Han.

The canonical dual frame of a wavelet frame.

*Appl. Comp. Harmonic Anal.* **12** (2002), no. 3, 269–285.

# The Frame Operator

## Proposition

Suppose that  $\Psi = \{\psi_1, \dots, \psi_L\} \subset L^2(\mathbb{R})$  generates a wavelet system which is a Bessel sequence. Let

$$\mathcal{D} = \{f \in L^2(\mathbb{R}) : \hat{f} \in L^\infty(\mathbb{R}) \text{ and } \text{supp } \hat{f} \subset [-R, -1/R] \cup [1/R, R] \text{ for } R > 1\}.$$

Then its frame operator  $S$  is given by

$$\widehat{Sf}(\xi) = \hat{f}(\xi) \sum_{l=1}^L \sum_{j \in \mathbb{Z}} |\hat{\psi}_l(2^j \xi)|^2 + \sum_{p \in \mathbb{Z}} \sum_{q \in 2\mathbb{Z}+1} \hat{f}(\xi + 2^{-p}q) t_q(2^p \xi)$$

for a.e.  $\xi \in \mathbb{R}$  and for all  $f \in \mathcal{D}$ , where

$$t_q(\xi) = \sum_{l=1}^L \sum_{j=0}^{\infty} \hat{\psi}_l(2^j \xi) \overline{\hat{\psi}_l(2^j(\xi + q))} \quad \text{for } q \in \mathbb{Z}.$$

This proposition is implicitly contained in the book of Hernández and Weiss, Proposition 7.1.19.

# Lifting

We are in some sense lifting the duality to the translation lattice  $P\mathbb{Z}$  and paring

$$\{D^j T_{Pk}(\psi), D^j T_{Pk}(T_1\psi), \dots, D^j T_{Pk}(T_{P-1}\psi)\}_{j,k \in \mathbb{Z}}$$

with

$$\{D^j T_{Pk}(\phi_0), D^j T_{Pk}(\phi_1), \dots, D^j T_{Pk}(\phi_{P-1})\}_{j,k \in \mathbb{Z}},$$

or, equivalently, lifting the duality to scale  $m$  and paring

$$\{D^j T_k(D^m \psi), D^j T_k(D^m T_1 \psi), \dots, D^j T_k(D^m T_{P-1} \psi)\}_{j,k \in \mathbb{Z}}$$

with

$$\{D^j T_k(\tilde{\phi}_0), D^j T_k(\tilde{\phi}_1), \dots, D^j T_k(\tilde{\phi}_{P-1})\}_{j,k \in \mathbb{Z}},$$

where  $\tilde{\phi}_i = D^m \phi_i$  for  $i \in \{0, 1, \dots, P-1\}$ . We note that the frame operator  $S$  remains the same during this lifting.